

Further Expanding the Metabolic Topography for Sustenance of Completeness Along with Working of Retina "-A Review

Kulvinder Kochar Kaur

Kulvinder Kaur Centre for Human Reproduction

Punjab, India

Abstract

Previously we reviewed the pivotal part of OS in the generation of NDD for instance ALS, AD, PD as well as alterations in brain energy metabolism particularly neuron-astrocytes energy metabolism in NDD inclusive of cross-cellular redox alignment associating i) astrocytic, ii) neuronal, as well as iii) microglial antioxidative event networks particularly how lactate shuttle significantly for astrocytes, whereas neurons reliability on glucose, besides part in ACh generation. Here we further illustrate the manner energy metabolism in retina is akin to neurons. Neurons possess considerably greater metabolic needs that are practically solely met by glucose provision from the bloodstream. Glucose is utilized in complicated nature of metabolic crosstalk amongst neurons as well as glia cells, displayed by the astrocyte-neuron lactate shuttle (ANLS) posit. The neural retina faces corresponding energy requirements in contrast to the rest of the brain, with greater extra anabolic demand to embrace persistent replenishment of photoreceptor outer segments. This needs is taken care of by an astonishing differences of the ANLS in that photoreceptors are the central part of a metabolic topography, with utilization of glucose in addition to supplying encompassing cells with metabolic intermediates. Here we describe recent evidence on the manner neurons, in particular photoreceptors, take care of their energy as well as biogenerational needs by constituting a metabolic panorama of , cells mutual reliability. supplied provision

Keywords:metabolic panorama; astrocyte-neuron lactate shuttle (ANLS);retina

INTRODUCING GLUCOSE METABOLISM

Despite cells meet several variable working, their working capability eventually has a fulcrum on the cell's capability of producing energy for survival as well as cell multiplication . The energy currency of cells is ATP, enabling thermodynamically unattractive biological reactions. Based on the accessibility of facilities , unicellular organisms possess the capacity of switching their metabolism amongst using nutrients to generate energy (catabolism) in addition to consuming such energy to generate the producing blocks of a new cell (anabolism). Multicellular organisms experience analogous need-guided switching in metabolism.


These are specifically pronounced in tissues that are multiplying however, further takes place in postmitotic cells, that possess the capability of switching toward anabolic states [1,2].

Catabolic reactions imperatively comprise of degrading molecules into smaller constituents that get oxidized to either yield energy or produce generating blocks for

anabolic reactions. The, most widely used carbon source is glucose, which is eventually catabolized to gain ATP, however, further possess the capacity of yielding pivotal precursors for biosynthetic reactions. There are two main routes of glucose metabolism: Either, i) glucose feeds into glycolysis, where it is ii) oxidized and becomes pyruvate, iii) yielding ATP as well as metabolic intermediates; iv) or alternatively, glucose get shunted from glycolysis into the pentose phosphate pathway for nucleic acid generation in addition to the v) generation of NADPH (**Figure 1**). [3].

Courtesy ref no.3-Metabolic pathways of glucose metabolism. Delineation of pivotal steps in glucose metabolism. Subsequent to gaining entry into the the cell, transformation of glucose takes place at the beginning by HK into G6P, utilizing ATP. G6P possesses the capability of branching off into the PPP pathway, generating R5P as well as creating large amounts of NADPH in the form of a byproduct.

Corresponding Author: Kulvinder Kochar Kaur
Kulvinder Kaur Centre for Human Reproduction
Punjab, India



Legend for Fig1

R5P is a precursor for generating nucleic acids. NADPH formed by the PPP possesses the capability of getting utilized to forage ROS through glutathione reduction, transforming H_2O_2 into H_2O . G6P is further metabolized into F6P in addition to, followed by into FBP. FBP is metabolized into one molecule of DHAP along with another of GADP. i) DHAP possesses the capacity of branching off into lipid biogenesis where it is a ii) precursor for phospholipids as well as iii) triacylglycerol. iv) Otherwise, DHAP is converted into v) GADP in addition to both a) GADP molecules are metabolized into b) 3-PG, c) creating ATP in the process. 3-PG possesses the capacity of getting utilized for i) serine biogenesis, which possesses the capacity of aiding in generation of ii) sphingolipids, along with iii) glycine as well as iv) cysteine. 3-PG goes on to be metabolized into v) PEP as well as subsequently into vi) pyruvate by PK activity. Pyruvate possesses the capacity of: working in the form of a) a substrate for alanine biogenesis; getting converted to lactate by LDH (referred to as anaerobic respiration or aerobic glycolysis); b) gaining entry into a mitochondrion where c) PDH metabolizes it into Acetyl-CoA. d) Acetyl-CoA subsequently gaining entry into the TCA cycle, where it is metabolized in a cascade of stepwise patterns reactions resulting in i) citrate, ii) α -KG, iii) succinate, iv) fumarate, v) malate, vi) oxaloacetate which cycles back to citrate. Certain of such reactions generate byproducts such as NADH and FADH_2 . NADH in addition to FADH_2 are subsequently utilized for OXPHOS, creating ATP in the inner mitochondrial membrane matrix.

Abbreviations: α -KG, α -ketoglutarate; 3-PG, 3-phosphoglycerate; ADP, adenosine diphosphate; ATP, adenosine triphosphate; CoA, coenzyme A; dATP, deoxyadenosine triphosphate; dCTP, deoxycytidine triphosphate; dGTP, deoxyguanosine triphosphate; DHAP, dihydroxyacetone phosphate; dTTP, deoxythymidine

triphosphate; ETC, electron transport chain; F6P, Fructose-6-phosphate; FADH, flavin adenine dinucleotide, reduced; FBP, fructose 1,6-bisphosphate; G6P, glucose-6-phosphate; GADP, glyceraldehyde-3-phosphate; GTP, guanosine triphosphate; HK, hexokinase; LDH, lactate dehydrogenase; NAD, nicotinamide adenine dinucleotide; NADH, nicotinamide adenine dinucleotide, reduced; NADP, nicotinamide adenine dinucleotide phosphate; NADPH, nicotinamide adenine dinucleotide phosphate, reduced; OXPHOS, oxidative phosphorylation; PEP, phosphoenolpyruvate; PDH, pyruvate dehydrogenase; PK, pyruvate kinase; PPP, pentose phosphate pathway; TCA, tricarboxylic acid.

Glycolysis is performed by a set of cytoplasmic enzymes ultimately yielding pyruvate, catalyzing the phosphorylation of two ADP to ATP as well as causing the reduction of two NAD^+ to NADH (Figure 1). In the presence of oxygen, pyruvate is oxidized to Acetyl-CoA in addition to enters the mitochondrial tricarboxylic acid (TCA) cycle (alias Krebs cycle or citric acid cycle). In the TCA cycle there are a series of oxidative reactions resulting in the generation of reductive power as in the form of NADH along with FADH^+ (Figure 1). The electron transport chain (ETC) in the membrane of the mitochondria then makes utility of this reductive power to form ATP in an event referred to as oxidative phosphorylation (OXPHOS). This mirrors the maximum profit constituents of energy metabolism owing to one glucose molecule forms as high as 36 molecules of ATP (Figure 1).

In case of restricted, oxygen accessibility is cells switching pyruvate away from the TCA and OXPHOS by reducing it to lactate in an event process referred to as anaerobic respiration. In the context of oxygen-restricted situations, NADH does not possess the capacity of being reduced to NAD^+ . Owing to need for

NAD^+ is existent even now in the form of an electron acceptor, pyruvate gets reduced to lactate, using NADH along with renewal of the pool of NAD^+ in the event. In the context of particular episodes, anaerobic respiration further possesses the capability of taking place when accessibility of oxygen enrichment is present, the manner detailed subsequently.

Apart from representing the backbone of energy metabolism, glycolysis as well as the TCA cycle further yield imperative precursors for biogenesis pathways. Plethora of glycolytic intermediates helps toward i) amino acid biogenesis, ii) nucleotide biogenesis, iii) complicated carbohydrate biogenesis, in addition to lipid biogenesis (**Figure 1**). The initial step of glycolysis transforms glucose into Glucose-6-phosphate (G6P) by hexokinase (HK) enzymes. G6P possesses the capacity of i) either getting ultimately metabolized to pyruvate, ii) or get shuttled to the pentose phosphate pathway (PPP), which yields the constructing blocks for nucleotides in addition to renewal of NADPH (**Figure 1**). The other noticeable biogenesis aids in glycolysis is generating precursors for phospholipid generation. The glycolytic intermediate dihydroxyacetone phosphate (DHAP) possess the capacity of getting oxidized to glycerol-3-phosphate, which is an essential precursor of diacylglycerophospholipids as well as glycerophospholipids, structural constituents of all biological membranes. Corresponding to glycolysis, the TCA cycle provides important precursors for lipid in addition to amino acid biogenesis (**Figure 1**).

Whereas the catabolic as well as anabolic parts of the

detailed pathways are imperative for all cells, variations working diverse metabolic needs are present. This is specifically correct for neurons.


Owing to their particular physiology in addition to working, neurons are energetically needy, along with depend practically solely on glucose for the energy as per their requirements [4,5]. The specific significance of neuronal energy metabolism arises from the slopy energy requirements of neuronal signaling. Neurons work by transitory altering their membrane potential (MP), that in turn describes a high metabolic load on regaining resting potential via the activity of ATP-based ion pumps.

Subsequently, our concentration is on energy metabolism of neurons in the brain in addition to the retina, along with try to present an integral model of retinal energy metabolism.

2. BRAIN ENERGY METABOLISM

The brain is among the maximum metabolically active organs as well as it depends practically solely on glucose in the form of its energy in the form of its energy facility [5-8]. It contribute to approximately 20% of entire body oxygen full utilization in addition to total 25% of glucose utilization whereas constituting just roughly 2% of total body weight [9,10]. Brain working depends on the relationship amongst neurons, glia, along with vasculature to meet such substantial metabolic needs (**Figure 2**).

Courtesy ref no.3-Neuron-Astrocyte metabolic interdependence. Delineation of the metabolite flow



Legend for Fig2

between blood vessels, astrocytes as well as synaptic terminals of neurons. Glucose provided from the blood passes the vessel endothelium in addition to, enveloping astrocyte endfeet via GLUT1. Glucose subsequently enters

aerobic glycolysis, with marginal contribution toward the TCA through due to reduced PDH activity. Resulting pyruvate is practically solely transformed into lactate through via the activity of LDH5. LDH5 is constituted of

LDHA subunits that possess propensity guide drive the reaction toward consuming NADH along with creating lactate, refilling the NAD⁺ pool for increased glycolytic activity. Lactate is subsequently exported through via MCT1 as well as MCT4 in addition to it is taken up by neurons through MCT2. In the neuron, lactate is metabolized back into pyruvate through LDH1. LDH1 is basically constituted of LDHB subunits that possess predilection to guide the reaction toward consuming NAD⁺, along with producing pyruvate, top up replenishing the pool of available NADH. Pyruvate fuels the TCA cycle as well as OXPHOS, creating ATP essential for refilling pre-synaptic glutamate vesicles. subsequently Upon neuronal activation, liberation of glutamate is r into the synapse in addition to binds to post-synaptic mGLUR GPCRs, ultimately resulting in a Ca²⁺ influx, along with signal propagation. Free glutamate in the synaptic cleft is re-uptaken by adjacent astrocytes through EAATs that co-transport Na⁺ in addition to K⁺. Glutamate gains entry into the the glutamate-glutamine cycle and is metabolized into glutamine by GS. Glutamine is subsequently exported, along with taken up by neurons, where it is returned back into glutamate by GLS, along with is utilized to refill glutamate vesicles. Alternatively, glutamate is metabolized into α -KG that enters the TCA cycle as well as supplements ATP production. ATP is subsequently utilized by Na⁺/K⁺ ATPases to re-generate the ionic gradient, renewal of the available ADP pool to stimulate glycolysis. Abbreviations: α -KG, α -ketoglutarate; ADP, adenosine diphosphate; ATP, adenosine triphosphate; EAAT, excitatory amino acid transporter; GCPR, G-protein-coupled receptor; Glc, glucose; Gln, glutamine; GLS, glutaminase; Glu, glutamate; GLUT, glucose transporter; GS, glutamate synthase; Lac, lactate; LDH, lactate dehydrogenase; MCT, monocarboxylate transporter; mGLUR, metabotropic glutamate receptor; NAD, nicotinamide adenine dinucleotide; NADH, nicotinamide adenine dinucleotide, reduced; OXPHOS, oxidative phosphorylation; PDH, pyruvate dehydrogenase; PKM1/M2, pyruvate kinase isozyme M1/M2; Pyr, pyruvate; TCA, tricarboxylic acid. possess utilization of

Astrocytes portray the maximum enriched as well as maximum germane glia cells for brain energy metabolism[11,12]. They possess plethora of parts in sustenance of appropriate brain working, certain of which are directly associated with energy metabolism[8,13]. Astrocytes modulate the association amongst neuronal energy demands, along with circulating glucose by directly touching in addition to encompassing blood vessels with projections (endfeet), as well as controlling blood supply germane to neuronal needs [14,15], by allowing for vasomotor responses[16]. Astrocytes work in the

form of the major entry point of glucose into the brain through glucose transporters (GLUTs) expressed at both endothelium as well as astrocyte endfeet (**Figure 2**). Such working relationship generates a pivotal neurometabolic coupling that is the benchmark of brain energy metabolism, making glia to sense neuronal energy needs in addition to adjust the provision as per that accordingly [17,18].

Whereas both neurons, along with glia account for brain energy metabolism, neuronal needs are substantially greater in contrast to glia's [19-21]. The major energy expenditure of neurons is correlated with re-generating MPs as well as making synaptic transmission[19,21,22]. Therefore the action of ion pumps for instance the Na⁺/K⁺ ATPases contribute to the maximum of total energy spent in the entire brain[19,23]. This is correct apart from for re-generating membrane ion gradients subsequent to membrane depolarization, however further in reference to sustenance of baseline MP that compensates spike-associated ion switching [19]. It has been determined that sustenance of baseline MP is implicated in reference to intricate to 50% of the total energy utilization in the white matter[24], in addition to approximately a 1/4th in cortical neurons [25]. Astonishingly, membrane depolarization by itself does not need substantial levels of ATP, owing to action potentials (AP) possess the capacity of remaining for certain time without ATP consumption[22,29]. Conversely, appropriate Na⁺ as well as K⁺ gradients Na⁺/K⁺ ATPase subsequent to AP action employs an extensive quantity of ATP (as high as 8×10^8 molecules). [25-28]. Intriguingly, in synaptic transmission Na⁺/K⁺ ATPase action delineates just a miniscule segment of energy expenditure in contrast to the needs of the synaptic vesicle cycle [19,22,29].

Whereas neurons are basically oxidative in their metabolism, depending on OXPHOS activity to generate ATP in an action-based manner [30-33].

In reference to the rectification of the appropriate Na⁺ along with K⁺ gradients subsequent to an AP.

Intriguingly, in synaptic transmission Na⁺/K⁺ ATPase action-based fashion[30], (whose subunit constitution attracts catalysis of pyruvate into lactate[36-39]. astrocytes are substantially glycolytic [34,35]. (**Figure 2**). Numerous factors estimate astrocytes' metabolic profile: i) First, astrocytes possess an enhanced expression in addition to action of Phosphofructokinase (PFK), thereby facilitating glycolysis[11,34,36]. ii) Secondly, astrocytes express lactate dehydrogenase 5 (LDH5), an enzyme in reference to rectification of the appropriate Na⁺ along with K⁺ gradients subsequent to an AP consumes an extensive quantity of ATP (up to 8×10^8 molecules)

[11,25,36,37]. iii) Lastly, astrocytes possess diminished pyruvate dehydrogenase (PDH) actions, that points that any escalation from basal pyruvate quantities shifts into LDH5- modulated transformation to lactate rather than gaining entry into the TCA cycle as well as OXPHOS actions [36,40,41]. (**Figure 2**). On the other hand, neuronal glycolysis quantities are negatively impacted by diminished generation in addition to actions of PFK[34], an incapacity of neuronal pyruvate kinase (PKM1) to regulate glycolytic flux, along with lastly, by expressing LDH1, a LDH isozyme whose subunit constitution kinetically attracts catalysis of lactate to pyruvate[34,42]. The amelioration of glycolysis in neurons allows glucose to be shifted to the PPP in reference to offsetting ROS production. As per that upregulating glycolysis in neurons is met with escalated oxidative stress (OS) as well as cell death[34,43,44].

The variation amongst neurons in addition to astrocytes is briefly detailed here is at the basis of their relationship as well as importantly shapes brain energy metabolism. This mutual reliability is formulated in the astrocyte-neuron lactate shuttle (ANLS) posit, one of the maximum broadly agreed upon models detailing brain energy metabolism at the cellular level [8,45-47]. Behind such model is the notion that despite glucose is the exclusive energy facility of the brain, glucose- obtained metabolites for instance lactate possesses the capacity of themselves working in the form of an alternate substrate for energy generation.

The basic postulate of the ANLS is neuronal activation that stimulates an escalation of astrocyte glucose uptake, that is subsequently basically transformed into lactate as well as shunted to neurons where it possesses the capability of in addition to getting used in the form of a substrate for oxidative energy metabolism[8,45] (**Figure 2**). In the ANLS original study, Pellerin and Magistretti[45], illustrated that glucose uptake in addition to sequential lactate synthesis in cultured astrocytes is triggered by the excitatory neurotransmitter glutamate[45].


Glutamate liberated into the synaptic cleft is picked up by excitatory amino acid transporters (EAATs), which are present in glial lamellar processes sheeting the post-, along with pre-synaptic elements[17]. The magnitude of glutamate import works in the form of a neuronal activity measure as well as glutamate uptake by EAATs results in a simultaneous influx of Na^+ in addition to efflux of K^+ . This triggers Na^+/K^+ ATPase activity thereby decreasing local ATP concentration, which in turn promotes glycolysis in astrocytes[35,36,48,49] (**Figure 2**). Similar to neurons, astrocytes in differentially fashion express a pyruvate kinase isoform (PKM2) that, compared to the neuronal isoform, possesses the capability of controlling glycolytic flux in reaction to escalated energy needs[36,50].

Astrocytic glycolysis feeds into lactate generation which is liberated via committed monocarboxylate transporters (MCTs) [51]. Neurons subsequently pick up extracellular lactate through neuron particular MCT where it is subsequently transformed into pyruvate by LDH1. Pyruvate subsequently gains entry into the TCA cycle, eventually generating amongst 14 as well as 17 molecules of ATP via OXPHOS per molecule of lactate [45,51,52] (**Figure 2**).

The substantial quantity of the earlier work behind the ANLS in addition to in the decades subsequently to its emergence is based on *in vitro* cell models of neurons, along with astrocytes. Debates in reference to the ANLS mostly encompass the query the manner easily such *in vitro* results are applicable to the situation in the intact brain[5,53]. Nonetheless, due to present work starts to concentrate on tissue culture as well as *in vivo* models, most of such queries are achieving replies [48,49,54-64]. Whereas the ANLS has been postulated to exposition the metabolic panorama for the whole central nervous system (CNS), the classical ANLS model does not possess the capacity of fully detailing the found metabolic crosstalk in the retina, being a part of the CNS. In the following section, we will amalgamate in addition to theoretically frame what is known in reference to the retinal energy metabolism.

3. RETINAL STRUCTURE ALONG WITH WORKING

In humans, vision is the primary sense as well as a substantial part of the neocortex is devoted to the processing of visual information. The initial step in reference to visual discrimination is light estimation in addition to preprocessing by the neural retina (**Figure 3A**). This conventionally layered tissue possesses three nuclear layers i) (ONL, INL, GCL), along with ii) two synaptic layers (OPL as well as IPL) possessing all of five variable neuronal in addition to one glial cell kind (**Figure 3B**). i) The ONL possesses a) rod in addition to b) cone photoreceptors, whereas ii) cell bodies of bipolar, a) amacrine as well as b) horizontal cells, c) in addition to Müller glia constituted the INL. iii) The third layer is the GCL, which possesses a) ganglion cells in addition to b) displaced amacrine cells. All of such cells account for modulating as well as processing signals formed by photoreceptors in reaction to light[65,66] (**Figure 3B**).



Legend for Fig3

Courtesy ref no.3-Vertebrate retina structure as well as cellular constitution. (A) Sagittal Delineation of the human eye. Light enters the eye through the lens in addition to gets projected on to the retina. (B) conventionally multilayered structure of the classical vertebrate retina. Light transverses the whole retina until it reaches the photosensible PhRs at the distal ONL, nearby with RPE projections that are neighbouring to the choroid plexus where choroidal vasculature supplies nutrients to the outer retina. In the OPL, PhR terminals touch projections from HC in addition to BC, whose cell body reside in the INL. Projections from BC touch those from the GC along with AC in the IPL. GC cell bodies positioned in the GCL as well as their projections bundle in the NFL to form the optic nerve. MGCs span all of such layers along with embrace the cells they touch. (C) Rod as well as cone PhRs cell structure. PhRs have an inner in addition to outer segment. PhR OSs house membranous disks that house the photosensitive pigment implicated for light detection. The IS consists of the rest of the cell body and houses a mitochondrial cluster at the base of the OS. Abbreviations: AC, amacrine cell; BC, bipolar cell; GC, ganglion cell; HC, horizontal cell; INL, inner nuclear layer; IPL, inner plexiform layer; IS, inner segment; MGC, Müller glia cell; NFL, nerve fiber layer; ONL, outer nuclear layer; OPL, outer plexiform layer; OS, outer segment; PhR, photoreceptor; RPE, retinal pigment epithelium.

Photoreceptors portray the light-sensitive neurons, which are implicated for transforming light into a biological signal in an event referred to as phototransduction. Both rod as well as cone photoreceptors possess an outer segment (OS) that possess membranous disks loaded with light-sensitive photopigments (Figure 3C). The photoreceptor outer segment is directly touching the

retinal pigment epithelium (RPE), implicated in plethora of embracing working in addition to the apical part of the RPE is touching the choroidal vasculature [67] (Figure 3B). This communication mirrors the photoreceptors' major facility of imperative nutrients, generating a major metabolite supply-line in reference to photoreceptors in the form of part of the Blood-Retina Barrier (BRB) [68].

3.1 Energy Metabolism of Retina


The specification of photoreceptor form along with working are at the root of their considerably greater energy demands [69,70]. Warburg [71], initially revealed the quantities of oxygen utilization of the retina, which was the maximum of any tissue of the body that he assessed, analogous just to swiftly proliferating cancer cells [71,72]. Warburg was pioneer for detailing the eponymic Warburg action in the form of a metabolic actuate to glycolysis in the existence of oxygen (aerobic glycolysis) illustrated by tumor cells. Apart from cancer cells, the Warburg action is further **observed** in the retina [73,74]. This is astonishing, due to compared with cancer cells, retinal neurons are postmitotic.

In addition to the greater catabolic requirement of photoreceptors, they illustrate skyrocketing anabolic requirements. This is in view of the curious circumstances of outer segment renewal, where photoreceptors are persistently shedding the outer most region of the outer segments, equivalent to approximately 10% per day [75,76]. Sequentially, new disks need to be persistently added at the base of the outer segment, needing a persistent provision of a plethora of large quantities complex macromolecules for instance lipids as well as proteins to in reference to sustenance of appropriate working and OS length [77-80]. The retina, analogous to

the rest of the CNS, possesses a predilection of utilization of glucose in the form of its main energy substrate. Glucose provision of the retina is basically modulated by glucose transporters[81,82], that are expressed in

vasculature, RPE cells, PhRs as well as in MGCs[70,83,84] (Figure 4).

4



Legend for Fig4

Courtesy ref no.3- Metabolic panorama of the vertebrate outer retina. Delineation of acknowledged metabolic crosstalk of cells amongst the outer retina. Glucose from the choroidal blood supply passes the endothelium as well as gains entry into the RPE through GLUT1. In the RPE, glucose flows via the cell un-metabolized, hence arriving at the basal membrane in the neighbouring PhR OS. Glucose escapes the RPE through GLUT1 into the PhR layer. PhRs pick up glucose via GLUT1 in addition to utilize it to fuel aerobic glycolysis.

Aerobic glycolysis in PhRs fuels the PPP, producing nucleic acids, along with offsetting ROS production from mitochondrial activity, as well as creates DHAP which is utilized to produce phospholipids imperative to produce OS disks hence guiding OS growth in addition to PhR working. Lactate from aerobic glycolysis is transported from PhRs to the extracellular space where it is picked up by other cells for instance MGC as well as RPE, fueling TCA cycle in addition to OXPHOS in such cells. RPE cells pick up lactate, along with convert it into pyruvate, that gains entry into the TCA. Such LDH activity leads to removal of accessible NAD⁺, efficaciously hampering glycolysis, accounts for glucose to arrive at GLUTs in the basal RPE membrane. Extension of OS tips externalize PS as well as touch RPE projections as well as following phagocytosis result in Akt activation, that further downregulates glycolysis in such cells. PhR OS phagocytosis in the RPE result in an escalation of accessible fatty acids (FAs) from breaking down the membranous lipid disks. Such FAs undergo

β -oxidation creating ketone bodies β -hydroxybutyrate, that possess the capacity of being also metabolized in addition to, utilized in the TCA cycle, supplementing RPE mitochondrial activity. Transportation of both FAs along with ketone bodies takes place to the PhR layer where they are picked up by as well as utilized to supplement TCA cycle in addition to OXPHOS. An existence of reciprocal malate-succinate shuttle amongst PhRs, along with RPE cells occurs that exchanges reducing power amongst such cells. In PhR, reverse SDH activity, owing to hypoxia-stimulated diminishing of COXIV, generates succinate that is exported to RPE cells, where it is oxidized in the TCA cycle in addition to produces an excess of malate. Malate is subsequently exported to the PhR layer, that gains entry into the TCA cycle, along with fuels succinate generation. Rods efficaciously facilitate cone survival as well as working by liberating RdCVF, that binds in addition to creating a complex with BSG1, along with GLUT1 in the cone membrane, facilitating glucose supply to such cells. Abbreviations: ADP, adenosine diphosphate; Akt, protein kinase B; ATP, adenosine triphosphate; BSG1, basigin 1; COXIV, cytochrome c oxidase subunit 4; DHAP, dihydroxyacetone phosphate; Glc, glucose; GLUT, glucose transporter; Lac, lactate; LDH, lactate dehydrogenase; Mal, malate; MCT, monocarboxylate transporter; MGC, Müller glia cell; NAD, nicotinamide adenine dinucleotide; NADH, nicotinamide adenine dinucleotide, reduced; OS, outer segment; OXPHOS, oxidative phosphorylation; PDH, pyruvate dehydrogenase; PhR, photoreceptor; PPP,

pentose phosphate pathway; PS, phosphatidylserine; Pyr, pyruvate; RdCVF, rod-derived cone viability factor; ROS, reactive oxygen species; RPE, retinal pigment epithelium; SDH, succinate dehydrogenase; Suc, succinate; TCA, tricarboxylic acid.

GLUTs mirror facilitative transporters, along with as such glucose observes a concentration gradient [85,86], provision of glucose from the blood leaves the choroidal endothelium as well as diffuses via RPE cells into the photoreceptor layer where photoreceptors pick it up in addition to metabolize it with ease. This facilitated diffusion is believed to be by GLUT1 [70,87].

Parallel to other neurons, the substantial segment of ATP expenditure amongst photoreceptors is an active ion transport along with repolarization of MP. ATP utilization in photoreceptors are activity based as well as considerably different in darkness in addition to in light [88,89]. The glutamate based dark current is energetically costly as well as needs a considerable quantity of ATP in reference to sustenance. In mice, rod photoreceptor consumes approximately four fold greater ATP in darkness in contrast to in light (about 10^8 ATP s^{-1}) [88]. Surplus, Ca^{2+} as well as Na^+ is actively transported out of the cell via ATPase pumps in addition to carries a meaningful energy expenditure.

The combined expenditure for such efflux is practically $9 \times 10^7 \text{ ATP s}^{-1}$. Intriguingly, energy profiles for rods along with cones vary quite considerably. Generally, cones possess greater energy demands in contrast to rods, plausibly owing to variations in energy needs of the dark current, phototransduction, as well as synaptic liberation [88,90]. The manner in most cells, a critical role of glucose in the retina is enabling ATP production to meet the cell's energy demands [73,83,91]. Both glycolysis as well as OXPHOS produce ATP, despite OXPHOS provided are substantially greater. It is by now well generated that photoreceptors generate lactate in the existence of oxygen [92,93]. The manner aforementioned, aerobic glycolysis further commonly labelled in the form of the Warburg effect, which is a common property of swiftly proliferating cancer cells in addition to neural stem cells [94]. Why should energy-needy cells for instance photoreceptors withdraw from utilization of OXPHOS, the maximum profit attaining energy generating pathway? Minimal exposition might be existent partly done by in the robust requirement of anabolic intermediates, yielded by glycolysis [87,95]. Despite, photoreceptors do not proliferate, they nonetheless possess a substantial anabolic need owing to the persistent renewal of their outer segments [73,75,76]. Despite, the craze over retinal energy metabolism has a practically century old-

history beginning with Warburg, greater recent research presented now provides the fundamentals in reference to the greater incorporating perspectives of retinal energy metabolism. The concentration has shifted away from exclusive photoreceptors however on the metabolic network of adjacent cells in the outer retina forming a metabolic panorama (Figure 4).

The initial thing to be taken into account is the manner glucose gains entry into the retina. Glucose is yielded by blood circulation along with needs to traverse the blood retina barrier (BRB). It is shunted via the choroidal endothelium into neighbouring RPE cells, prior to arriving at the outer retina. RPE shuttles maximum glucose via GLUT1 to the photoreceptor layer, whereas depending on lactate provided by photoreceptors for its own metabolism [82,93,96]; (Figure 4). RPE cells ensure least glucose utilization along with results in maximization of glucose efflux by utilization of photoreceptor-obtained lactate in the form of their metabolic substrate. Simultaneously, lactate is a negative feedback regulator of glycolysis that impedes RPE glucose utilization [93] (Figure 4). Such intricate metabolic co-reliability is not restricted to catabolism, but extends to anabolism. RPE cells export malate for photoreceptors to refill succinate that is exported back to RPE cells. Such reaction makes RPE cells to generate reductive power in the form of NADPH regenerating malate in the event, hence closing the cycle [97] (Figure 4). Malate possesses the capability of further being converted to pyruvate, yielding additional NADPH in the event [98] (Figures 1, 4). RPE cells are further acknowledged to depend on β -oxidation of fatty acids (FAs), resulting from phagocytosed disks of shed photoreceptor OS, to generate energy. The resulting KBs supplement the TCA cycle, providing ATP in addition to metabolic intermediates, as well as hence compensate for the downregulation of glycolysis [99-101] (Figure 4). Moreover, the RPE has been illustrated to provide photoreceptors with FAs for as well as ketone bodies (KBs), which possess the capacity of getting metabolized in the mitochondria in addition to supplement TCA cycle activity, along with ATP generation [99,101,102] (Figure 4). One additional cornerstone of the mutual reliability amongst photoreceptor outer segments in addition to the RPE is the recent observation that touches amongst phosphatidylserine on outer segment tips as well as RPE receptors activates the intracellular signaling Akt pathway. This efficaciously associates phagocytosis with glucose transportation to photoreceptors for new outer segment generation [103] (Figure 4).

Collectively, such studies illustrate an intricate metabolic crosstalk amongst the RPE as well as the outer retina in reference to the exchange of apart from glucose in addition

to, lactate, however further metabolites implicated in anabolism. The involvement of aerobic glycolysis in biogenesis along with maintenance of outer segment has been illustrated further in rod photoreceptors, by disturbing pivotal enzymes mediating aerobic glycolysis [104,105] .

In the metabolic topography there are apart from reciprocal crosstalk amongst the RPE as well as photoreceptors, however further amongst rods in addition to cones. The mutual reliability of photoreceptors has been acknowledged for substantial time period by the confusing 'onlooker actions' where genetically jeopardized rods result in the death of non jeopardized cones[106] . Such action is posited to be in view of the lack of the rod-derived cone viability factor(RdCVF) that occur secondary to rod degeneration. The splice variant of the *Nxn1* genes encodes RdCVF, where binding takes place with Basigin-1 (BSG1) expressed on the cell surface of cones. BSG1 creates a complex with GLUT1, stimulating glucose uptake upon RdCVF binding along with sequentially aerobic glycolysis[107,108] ; **Figure 4**). The lack of such crosstalk causes metabolic starvation of cones.

In the posited metabolic panorama setting, photoreceptors are set in the form of energy relay nodes which utilize glucose regarding attaining their anabolic as well as catabolic requirements. Simultaneously , they yield encompassing cells with metabolic intermediates (**Figure 4**). This is a curious difference on the ANLS in other areas of the CNS. We watched the manner the ANLS delineates a metabolic exchange amongst astrocytes as well as neurons where astrocytic lactate from aerobic glycolysis assists in fuel neuronal actions. Amongst the retinal metabolic panorama , photoreceptors pick on an analogous part to astrocytes in the remaining aspects of the CNS by provision of lactate to encompassing cells. Furthermore, both astrocytes in the CNS, along with photoreceptors in the retina are the first neural cells to touch glucose in their respective system. Such fascinating "reverting of part" might differ substantially in the form of adaption to the distinct micromilieu of the retina as well as the extraordinary anabolic requirements of postmitotic photoreceptors. The massive needs of glycolytic intermediates for anabolism makes photoreceptors act in the form of glucose sink (alias any tissue, organ, or mechanism in the body that absorbs, utilizes, or stores glucose) from the bloodstream that encompasses cells have created resilience to by utilizing by-products to embrace their own metabolic requirements [93].

Such model points to a close in addition to mutual reliability on cell milieu , yet there are certain intriguing angles that await further clarification. Among them is the metabolic differentiation amongst rod along with cone photoreceptors as well as utility - based metabolic requirements. Another curious extraordinary query is the manner photoreceptors tackle the lesser ATP- provision yielding aerobic glycolysis to embrace lone oxidative metabolism. Additionally, if this holds true then what is the need for photoreceptors possessing extensively big , intricately packed mitochondrial clusters neighbouring the OS[109] ? Whereas the present retinal metabolic panorama specifies plethora of crosstalks amongst the retina from a bioenergetics point-of-view, it is substantially incomplete as well as plethora of the proposed crosstalks would benefit from additional robustness in addition to autonomous confirmation. Additionally, practically all of the embracing outcomes is from murine models that depend mainly on scotopic (nocturnal) vision in addition to whose retina is comprised of only practically 2% cones. The rest of the data is from *in vitro* cell modeling of the retina. Whereas both such strategies have verified to be meaningful , there is a substantial requirement for further *in vivo* outcomes avoiding greater artificial *in vitro* methodologies. This is specifically correct for knowledge on cone photoreceptors, that modulate high-acuity color vision on that humans depend on.

3.2 Retinal Metabolism In Disease

Therefore here we have detailed recent evidence that glucose metabolism is pivotal to sustenance of brain in addition to retina homeostasis. Acknowledged the robust reliability of cells on glucose metabolism, it comes is not astonishing that metabolism is frequently influenced in neural diseases, be whether it acts in the form of the etiological factor or secondary as the results of metabolic disturbances. The retina is akin to this as well as plethora of retinal diseases have been associated with metabolism. A reason are the pathomechanisms of outer retinal dystrophies, for instance of age correlated - macular degeneration (AMD) in addition to, retinitis pigmentosa (RP). In such group of diseases, there is a growing acknowledgement that metabolic alterations might underlie as their pathophysiology [80,87,110] .

Age- correlated macular degeneration delineate's a retinal disease that is the maximum frequent etiological factor of blindness in elderly worldwide[111] . Whereas the mechanistic modes behind AMD continues to be uncharted , plethora of recent studies suggest oxidative injury [112-114]

as well as mitochondrial impairment in the form of pivotal


steps in AMD pathophysiology[80,87,115-117]. The idea of a retinal metabolic panorama possesses the capability of aiding in associating accrual of mitochondrial DNA injury in the RPE to macular degeneration. RPE cells with jeopardized mitochondrial working might switch toward escalated glucose consumption in addition to hence imperil photoreceptor access to glucose.

On the other hand, alterations in photoreceptor energy supply will have implications on the RPE.

3.2B Retinitis pigmentosa (RP) portray 's the disease maximum tightly associated with alterations in the retinal metabolic panorama. The retinal metabolic panorama brought forward here was robustly guided by studies associated with RP[78,103,104,108]. RP portray 's a group of rare diseases where mainly rod- particular mutations are etiological factors for rod, followed by cone, degeneration [118]. The expositions for such onlooker actions" possess the capacity of being accredited to the absence of cone viability factor (RdCVF), which ultimately stimulates cone glucose uptake[108].

The manner the complicated nature of the retinal metabolic topography begin to unravel, we start to look at the manner energy homeostasis is pivotal to sustenance of an appropriate working system. Disturbances in the metabolic crosstalk possess the capacity of possessing important sequelae resulting in particular disease phenotypes.

3.2C Dry age-correlated macular degeneration (AMD) is the commonest etiological factor of central vision elimination among the elderly, however yet no curative therapy is present. While exudative AMD possesses the capability of getting managed with anti-vascular endothelial growth factor (VEGF) treatment, dry AMD- contributes to greater than 85% of cases-propagates slyly from drusen accrual to geographic atrophy (GA). Despite the recent U.S. FDA recommendations of pegcetacoplan as well as avacincaptad pegol delineate main benchmark, their therapeutic effects continue to be modest. Bhandari et al [119], described an incorporated review of the molecular in addition to cellular mechanistic modes underlying dry AMD, emphasizing pivotal pathogenic pathways i) implicating OS, ii) lipid decontrolling, iii) complement activation, iv) mitochondrial dysfunction, along with v) RPE- particular bisretinoid lipofuscin accrual. Additionally, they assessed mechanistic mouse models which replicate such pathological events as well as describe the manner each model contributes to getting insight in the disease. Lastly, they described present in addition to appearing therapeutic approaches-including complement hampering agents, visual cycle modulators, along with mitochondrial-protective approaches- in addition to outline future trajectories for translational research. Taken together, their review synthesizes mechanistic insights, disease models, along with therapeutic innovation to embrace the generation of targeted as well as stage- particular mediation for dry AMD.




Legend for Fig5

Courtesy ref no.119-Pathological hallmarks of dry age-related macular degeneration (AMD). Dry AMD progresses from early drusen formation and intracellular lipofuscin accumulation to RPE degeneration, photoreceptor loss, and choriocapillaris thinning, ultimately leading to geographic atrophy.

Courtesy ref no.119-Pathomechanistic processes in dry AMD. Bisretinoid lipofuscin accumulation, phototoxic stress, lipid dysregulation, and complement activation

converge to induce oxidative injury, lysosomal dysfunction, and chronic inflammation in the RPE. A. Normal retina structure showing photoreceptors, retinal pigment epithelium (RPE), Bruch's membrane (BrM), and choroid. B. In early dry AMD, bisretinoid lipofuscin (e.g., A2E) accumulates in RPE lysosomes, with concomitant drusen formation and photoreceptor stress. C. Bisretinoids such as A2E, generated via the visual cycle, accumulate in RPE lysosomes. Upon light exposure, A2E undergoes




Legend for Fig6

photooxidation and photodegradation, producing singlet oxygen and superoxide radicals, leading to the generation of reactive degradation products. These photooxidative processes further impair lysosomal function and proteasome activity, reduce antioxidant capacity, and initiate chronic inflammation in RPE cells. This cascade contributes to two major pathogenic axes: (1) Complement dysregulation, initiated by downregulation of CFH and upregulation of CFB and C5b-9, promoting RPE cell apoptosis and inflammatory amplification. (2) Generation of oxidation-specific epitopes (OSEs), including advanced glycation end products (AGEs), carboxyethylpyrrole (CEP), malondialdehyde (MDA), 4-hydroxynonenal (4-HNE), and oxidized phospholipids (oxPLs)—arising from both A2E photodegradation and lipid peroxidation. These OSEs stimulate innate and autoimmune responses, driving further complement activation (e.g., C3, CFB, C5

upregulation) and suppression of complement inhibitors (CD46, CD55, CD59). In parallel, RPE-derived lipoprotein particles accumulate in BrM due to impaired efflux (e.g., via ApoE/A1, ABCA1, SR-BI) and compromised CFH regulation, especially in the Y402H risk variant. This leads to the formation of large HDL-like deposits (HLDs) with proinflammatory properties that further exacerbate early AMD lesion formation, such as drusen. RPE, retinal pigment epithelium; POS, photoreceptor outer segment; A2E, N-retinylidene-N-retinylethanolamine; *atRAL*, all-trans-retinal; *atROL*, all-trans-retinol; *atRE*, all-*trans*-retinyl ester; 11*cis*ROL, 11-*cis*-retinol; 11*cis*RAL, 11-*cis*-retinal; BrM, Bruch's membrane; ROS, reactive oxygen species; RDHs, retinol dehydrogenase; LRAT, lecithin retinol acyl transferase; RPE65, retinal pigment epithelium-specific 65 kDa protein.

Courtesy ref no.119-Complement pathways and therapeutic



Legend for Fig7

targets in geographic atrophy (GA). FDA-approved inhibitors (pegcetacoplan and avacincaptad pegol) act at C3 and C5 to suppress complement-mediated retinal damage.

To meet the substantial bioenergetic demands of the photoreceptors, glucose as well as other nutrients have to pass the retinal pigment epithelium (RPE), (that delineates a polarised monolayer of cells) that are located as the intermediary amongst the outer retina in addition to the choroid, the fundamental vascular layer of the eye. Recent evaluations have revealed a metabolic ecosystem in the outer retina where the photoreceptors, along with RPE engage in a complicated exchange of i) sugars, ii) amino acids, as well as iii) other metabolites. Disturbances of such delicate metabolic harmony has been isolated in the i) aging retina, in addition to in ii) age-associated macular degeneration (AMD), the maximum frequent etiological factor of blindness in the West. Further frequently in the aging, along with diseased retina are escalated quantities of i) cytokines, ii) oxidative stress (OS), iii) advanced glycation end-products, iv) escalated GF signalling, along with v) biomechanical stress – all of which have been associated with metabolic decontrolling in non-retinal cell kinds as well as tissues. Hansman et al [120], detailed the part of such factors in i) retinal homeostasis, ii) aging, as well as iii) disease. Further Hansman et al [120], outlined their actions on i) glucose, ii) mitochondrial, iii) lipid, in addition to iv) amino acid metabolism in tissues along with v) cell kinds outside the retina, emphasizing the signalling pathways via which they induce such alterations. Lastly, they detail attractive vistas for future research evaluating the roles of such pathological disorders on retinal metabolism, plausibly yielding innovative therapeutic strategies to tackle age-correlated retinal diseases [120].

4. CONCLUSIONS ALONG WITH FUTURE DIRECTIONS

Previously we reviewed the pivotal part of OS in the generation of NDD for instance ALS, AD, PD as well as alterations in brain energy metabolism particularly neuron-astrocytes energy metabolism in NDD inclusive of cross-cellular redox alignment associating i) astrocytic, ii) neuronal, as well as iii) microglial antioxidative event networks particularly how lactate shuttle significantly for astrocytes, whereas neurons reliability on glucose, besides part in ACh generation [121-130].

The aging RPE as well as Bruch's membrane are plagued by a range of pathological situations, in addition to others common to aging tissues, that possesses the capability of affecting RPE metabolism. These are inclusive of i)

chronic, ii) nonresolving inflammation guided by i) DAMPs, ii) cellular senescence, iii) basal deposits, along with iv) drusen; v) oxidative stress (OS) aggravated by a) high OXPHOS, b) photoreceptor OS phagocytosis, c) lipofuscin, d) incoming solar radiation, as well as e) diminishing antioxidant capacity; f) accrual of AGEs in Bruch's membrane g) with escalated expression of their receptor in the RPE; in addition to h) decontrolled MMP activity in Bruch's membrane, plausibly influencing growth factor (GF) signalling pathways. Additionally, aging is correlated with escalating biomechanical stress placed upon the RPE by i) basal deposits, ii) drusen, along with a iii) stiffening of Bruch's membrane. iv) Moreover, pathologies for instance AMD possesses powerful associations with i) chronic inflammation, ii) OS, iii) GF signalling, in addition to iv) metabolic decontrolling. v) Such pathological features might converge to result in metabolic decontrolling in the RPE, jeopardizing the capability of such cells to perform the pivotal function of promoting metabolite exchange between the choroid and outer retina. Greater work requirement is present to illustrate entirely the complicated association amongst such disease states, factors for instance i) cytokines, ii) growth factors (GF's), iii) free radicals, as well as iv) RPE metabolism. Gene editing technology advancements have extensively escalated the effectiveness of generating animal models to study the manner single genes or pathways influence different pathologies [131].

Mice where specific genes encoding i) antioxidant, ii) metabolic, in addition to iii) inflammatory constituents have been knocked out have confirmed to be strong gadgets for getting insight of the etiological factor, along with propagation of AMD [132]. Furthermore, cell-particular promoters have been utilized to knock out genes particularly in photoreceptors [104,133-135] as well as RPE cells [136-138]. Such animal models have yielded i) a precious gadget for illustrating apart from the part of specific genes in retinal disease, however further in inferring their imperative working in particular retinal cell kinds. Acknowledged our incepting understanding of the association amongst GF's cytokines, as well as other factors with the pathogenic properties of aging in addition to disease, animal models for instance these pose precious facilities, where imperatively any gene possesses the capacity of being knocked out specifically in defined cell kinds or tissues, despite under temporal regulation. Nevertheless, whereas such animal models helps the parts of specific genes in retinal generation along with visual working to be estimated, discriminating the specific actions on metabolism in distinct retinal cell populations, as well as the whole retinal metabolic ecosystem *in vivo*

poses greater tough botherations. Our insight of the retinal metabolic panorama has had extension swiftly in the recent decade. Advances in mass spectrometry technology now allow the detailed examination of the 'metabolome', the implementation of which has recently been applied to bulk retina in addition to RPE metabolomics [139-141]. The incorporation of radiolabelled metabolite therapies of cultured RPE cells [142-145] as well as explanted (akin to the removal of living tissue from an organism (plant or animal) to be placed in a nutrient medium for tissue culture, or the surgical removal of a foreign body, such as a medical device or prosthesis) retinal tissue [93, 142, 145, 146] has yielded insight into the manner the retina utilizes nutrients, along with emphasized the manner pivotal metabolites get exchanged amongst particular cell kinds. Whereas the metabolic evaluation of *ex vivo* tissue in addition to cultured cells provide useful understanding, there are limitations intrinsic to the elimination of these cells from their native state. The evolution of the field of retina metabolism is switching towards *in vivo* evaluation of metabolites, aiding in the determination of real-time alterations in flux in reactions to genetic as well as chemical disturbances. The refinement of existing in addition to emergence of novel methodologies possessing the capacity of determining metabolite enrichment amongst living organisms is starting to convert both the manner retinal metabolism is evaluated, along with the quality, exactitude as well as scrupulousness of data. Magnetic resonance spectroscopy (MRS) is a methodology that combines the principles of nuclear magnetic resonance (NMR) with determination utilizing an MRI scan. This non-invasive methodology has been useful gadget in deriving a fundamental canvas of the metabolic panorama of tumours as well as the brain [147, 148]. Nonetheless, whereas it persists to be a robust gadget it is plagued presently from the limitations of lesser sensitivity [148] in addition to resolution [149]. Despite such factors are anticipated to undergo improvement with advancements in ultra-high magnetic field buttressing along with coil technology [150], the resolution is still inadequate to detect metabolic changes in particular areas of the retina, substantially lesser amongst personal retinal cells. Microdialysis combined with mass spectrometry is one extra robust metabolomic methodology, useful for monitoring alterations in neurotransmitters, neuromodulators, as well as other metabolites in brain extracellular in addition to cerebral spinal fluid (CSF) [151, 152].

Such methodology offers the benefit of assisting in persistent fluid sampling from live animals. Microdialysis of vitreous fluid neighbouring the retina has been utilized

to monitor glutamate liberation in reactions to ischemia in mice [153], as well as 17- β glucuronide in rats [154]. Despite such methodology does not possess the capacity of inferring metabolic variations amongst personal retinal cells, it is capable of yielding in-depth understanding into the manner it influences the development in addition to liberation of metabolites *in vivo*, along with the transportation of particular metabolites across the BRB. A vital improvement in mass spectrometry has been the generation of mass spectrometry imaging (MSI), which yields the spatially resolved distribution of metabolites in tissues. This has promoted the propagation of the methodology from bulk, lesser resolution multi-cell evaluation of tissues to proteomics as well as metabolomics at single cell resolution [155, 156]. Whereas plethora of MSI procedures are present, evaluation of biological samples generally yield implicates the ionisation of metabolites in tissue sections, that gets followed by determination by the mass spectrometer. The spatial resolution of MSI is persistently escalating, now possessing the capacity of greater than 2 μ m resolution, in addition to is of determining metabolites with enrichment in the femtomole to attomole range utilizing secondary ion mass spectrometry (SIMS) [155, 156]. MSI has been utilized to measure metabolites in porcine retinas [157], as well as retinal lipids in mice [158]. Astonishingly, apart from such studies, that are practically over 10 yrs old, the implementation of MSI in retina research has been restricted. Despite innovations such as gas cluster ion beams (GCIBs) have substantially escalated the sensitivity, present methodologies have considerably lesser ionisation effectiveness which differ amongst metabolites, leaving substantial chance for improvement [155]. With generation of such methodologies, it is anticipated that the distinct metabolism of the RPE as well as lesser total of the other cells in the retina would have properties determined at magnitude of single cell *in vivo* for total of the main metabolites in addition to pivotal pathways. Once combined with genetic knockouts, along with other animal models, such methodologies would yield a robust gadget to decipher the manner metabolic exchange in the retina is interfered with by factors implicated in retinal aging as well as disease for instance OS, a stiff Bruch's membrane, AGEs, in addition to particular GF's as well as cytokines. Improved technology should meet the examination of other meaningful metabolites further than glucose metabolism, OXPHOS, along with associated pathways for instance the PPP by isolating alterations in constituents for instance lipids or amino acids. Generating the association amongst the aging as well as disease-germane factors as well as metabolic alterations

would apart from aid in our getting insight of retinal pathology, however might display therapeutic targets for instance pivotal factors, metabolites, as well as particular signalling constituents. Thus, getting insight of such events is of pivotal significance once we switch towards superior clinical outcomes in aging in addition to disease.

REFERENCES

- Vander Heiden, M. G., Cantley, L. C., and Thompson, C. B. (2009). Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* 324,1029–1033. doi: 10.1126/science.1160809
- Lunt, S. Y., and Vander Heiden, M. G. (2011). Aerobic glycolysis: meeting the metabolic requirements of cell proliferation. *Annu. Rev. Cell Dev. Biol.* 27,441–464. doi: 10.1146/annurev-cellbio-092910-154237.
- Viegas FO and Neuhaus SCF(2021) A Metabolic Landscape for Maintaining Retina Integrity and Function. *Front. Mol. Neurosci.* 14:656000. doi: 10.3389/fnmol.2021.656000
- Ames, A. (2000). CNS energy metabolism as related to function. *Brain Res. Rev.* 34,42–68. doi: 10.1016/S0165-0173(00)00038-2
- Dienel, G. A. (2019). Brain glucose metabolism: integration of energetics with function. *Physiol. Rev.* 99, 949–1045. doi: 10.1152/physrev.00062.2017
- Ashrafi, G., and Ryan, T. A. (2017). Glucose metabolism in nerve terminals. *Curr. Opin. Neurobiol.* 45, 156–161. doi: 10.1016/j.conb.2017.03.007
- Howarth, C., Gleeson, P., and Attwell, D. (2012). Updated energy budgets for neural computation in the neocortex and cerebellum. *J. Cereb. Blood Flow*
- Metab.* 32, 1222–1232. doi: 10.1038/jcbfm.2012.35 Magistretti, P. J., and Allaman, I. (2015). A cellular perspective on brain energy
- metabolism and functional imaging. *Neuron* 86, 883–901. doi: 10.1016/j.neuron.2015.03.035
- Mink, J. W., Blumenshine, R. J., and Adams, D. B. (1981). Ratio of central nervous system to body metabolism in vertebrates: its constancy and functional basis. *Am. J. Physiol.* 241, R203–R212. doi: 10.1152/ajpregu.1981.241.3.R203
- Sokoloff, L. (1977). Relation between physiological function and energy metabolism in the central nervous system. *J. Neurochem.* 29, 13–26. doi: 10.1111/j.1471-4159.1977.tb03919.x
- Barros, L. F., Brown, A., and Swanson, R. A. (2018). Glia in brain energy metabolism: a perspective. *Glia* 66, 1134–1137. doi: 10.1002/glia.23316
- Jha, M. K., and Morrison, B. M. (2018). Glia-neuron energy metabolism in health and diseases: new insights into the role of nervous system metabolic
- transporters. *Exp. Neurol.* 309, 23–31. doi: 10.1016/j.expneurol.2018.07.009 Bélanger, M., Allaman, I., and Magistretti, P. J. (2011). Brain energy metabolism: focus on astrocyte-neuron metabolic cooperation. *Cell Metab.* 14, 724–738. doi: 10.1016/j.cmet.2011.08.016
- Oberheim, N. A., Wang, X., Goldman, S., and Nedergaard, M. (2006). Astrocytic complexity distinguishes the human brain. *Trends Neurosci.* 29, 547–553. doi: 10.1016/j.tins.2006.08.004
- Gordon, G. R. J., Choi, H. B., Rungta, R. L., Ellis-Davies, G. C. R., and MacVicar, B. A. (2008). Brain metabolism dictates the polarity of astrocyte control over
- arterioles. *Nature* 456, 745–749. doi: 10.1038/nature07525 Attwell, D., Buchan, A. M., Charpak, S., Lauritzen, M., MacVicar, B. A., and Newman, E. A. (2010). Glial and neuronal control of brain blood flow. *Nature* 468, 232–243. doi: 10.1038/nature09613
- Iadecola, C., and Nedergaard, M. (2007). Glial regulation of the cerebral microvasculature. *Nat. Neurosci.* 10, 1369–1376. doi: 10.1038/nn2003
- Mathiisen, T. M., Lehre, K. P., Danbolt, N. C., and Ottersen, O. P. (2010). The perivascular astroglial sheath provides a complete covering of the brain microvessels: an electron microscopic 3D reconstruction. *Glia* 58, 1094–1103. doi: 10.1002/glia.20990
- Alle, H., Roth, A., and Geiger, J. R. P. (2009). Energy-efficient action potentials in hippocampal mossy fibers. *Science* 325, 1405–1408. doi: 10.1126/science.1174331
- Attwell, D., and Laughlin, S. B. (2001). An energy budget for signaling in the grey matter of the brain. *J. Cereb. Blood Flow Metab.* 21, 1133–1145. doi: 10.1097/00004647-200110000-00001
- Nortley, R., and Attwell, D. (2017). Control of brain energy supply by astrocytes. *Curr. Opin. Neurobiol.* 47, 80–85. doi: 10.1016/j.conb.2017.09.012

23. Rangaraju, V., Calloway, N., and Ryan, T. A. (2014). Activity-driven local ATP synthesis is required for synaptic function. *Cell* 156, 825–835. doi: 10.1016/j.cell.2013.12.042
24. Harris, J. J., Jolivet, R., and Attwell, D. (2012). Synaptic energy use and supply. *Neuron* 75, 762–777. doi: 10.1016/j.neuron.2012.08.019
25. Harris, J. J., and Attwell, D. (2012). The energetics of CNS white matter. *J. Neurosci.* 32, 356–371. doi: 10.1523/JNEUROSCI.3430-11.2012
26. Lennie, P. (2003). The cost of cortical computation. *Curr. Biol.* 13, 493–497. doi:10.1016/S0960-9822(03)00135-0
27. Kole, M. H. P., Ilschner, S. U., Kampa, B. M., Williams, S. R., Ruben, P. C., and Stuart, G. J. (2008). Action potential generation requires a high sodium channel density in the axon initial segment. *Nat. Neurosci.* 11, 178–186. doi:10.1038/nn2040
29. Howarth, C., Gleeson, P., and Attwell, D. (2012). Updated energy budgets for neural computation in the neocortex and cerebellum. *J. Cereb. Blood Flow Metab.* 32, 1222–1232. doi: 10.1038/jcbfm.2012.35
30. Hallermann, S., Kock, C. P. J., de Stuart, G. J., and Kole, M. H. P. (2012). State and location dependence of action potential metabolic cost in cortical pyramidal neurons. *Nat. Neurosci.* 15, 1007–1014. doi: 10.1038/nn.3132
31. Yuan, Y., Huo, H., and Fang, T. (2018). Effects of metabolic energy on synaptic transmission and dendritic integration in pyramidal neurons. *Front. Comput. Neurosci.* 12:79. doi: 10.3389/fncom.2018.00079
32. Verstreken, P., Ly, C. V., Venken, K. J. T., Koh, T.-W., Zhou, Y., and Bellen, H. J. (2005). Synaptic mitochondria are critical for mobilization of reserve pool vesicles at *Drosophila* neuromuscular junctions. *Neuron* 47, 365–378. doi:10.1016/j.neuron.2005.06.018
33. Ly, C. V., and Verstreken, P. (2006). Mitochondria at the synapse. *Neuroscientist* 12, 291–299. doi: 10.1177/1073858406287661
34. Hall, C. N., Klein-Flügge, M. C., Howarth, C., and Attwell, D. (2012).
35. Oxidative phosphorylation, not glycolysis, powers presynaptic and postsynaptic mechanisms underlying brain information processing. *J. Neurosci.* 32, 8940–8951. doi: 10.1523/JNEUROSCI.0026-12.2012
36. Lin, A.-L., Fox, P. T., Hardies, J., Duong, T. Q., and Gao, J.-H. (2010).
37. Nonlinear coupling between cerebral blood flow, oxygen consumption, and ATP production in human visual cortex. *Proc. Natl. Acad. Sci. U.S.A.* 107, 8446–8451. doi: 10.1073/pnas.0909711107
38. Herrero-Mendez, A., Almeida, A., Fernández, E., Maestre, C., Moncada, S., and Bolaños, J. P. (2009). The bioenergetic and antioxidant status of neurons is controlled by continuous degradation of a key glycolytic enzyme by APC/CCdh1. *Nat. Cell Biol.* 11, 747–752. doi: 10.1038/ncb1881
39. Bittner, C. X., Valdebenito, R., Ruminot, I., Loaiza, A., Larenas, V., Sotelo-Hitschfeld, T., et al. (2011). Fast and reversible stimulation of astrocytic glycolysis by K⁺ and a delayed and persistent effect of glutamate. *J. Neurosci.* 31, 4709–4713. doi: 10.1523/JNEUROSCI.5311-10.2011
40. Zhang, Y., Chen, K., Sloan, S. A., Bennett, M. L., Scholze, A. R., O’Keeffe, S., et al. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J. Neurosci.* 34, 11929–11947. doi: 10.1523/JNEUROSCI.1860-14.2014
41. O’Brien, J., Kla, K. M., Hopkins, I. B., Malecki, E. A., and McKenna, M. C. (2007). Kinetic parameters and lactate dehydrogenase isozyme activities support possible lactate utilization by neurons. *Neurochem. Res.* 32, 597–607. doi: 10.1007/s11064-006-9132-9
42. Mason, S. (2017). Lactate shuttles in neuroenergetics-homeostasis, allostasis and beyond. *Front. Neurosci.* 11:43. doi: 10.3389/fnins.2017.00043
43. Duka, T., Collins, Z., Anderson, S. M., Raghanti, M. A., Ely, J. J., Hof, P. R., et al. (2017). Divergent lactate dehydrogenase isoenzyme profile in cellular compartments of primate forebrain structures. *Mol. Cell. Neurosci.* 82, 137–142. doi: 10.1016/j.mcn.2017.04.007
44. Halim, N. D., Mcfate, T., Mohyeldin, A., Okagaki, P., Korotchikina, L. G., Patel, M. S., et al. (2010). Phosphorylation status of pyruvate dehydrogenase distinguishes metabolic phenotypes of cultured rat brain astrocytes and neurons. *Glia* 58, 1168–1176. doi: 10.1002/glia.20996

45. Hirrlinger, J., and Dringen, R. (2010). The cytosolic redox state of astrocytes: maintenance, regulation and functional implications for metabolite trafficking. *Brain Res. Rev.* 63, 177–188. doi: 10.1016/j.brainresrev.2009.10.003
46. Ward, P. S., and Thompson, C. B. (2012). Metabolic reprogramming: a cancer hallmark even warburg did not anticipate. *Cancer Cell* 21, 297–308. doi: 10.1016/j.ccr.2012.02.014
47. Gavillet, M., Allaman, I., and Magistretti, P. J. (2008). Modulation of astrocytic metabolic phenotype by proinflammatory cytokines. *Glia* 56, 975–989. doi: 10.1002/glia.20671
48. Bolaños, J. P., Almeida, A., and Moncada, S. (2010). Glycolysis: a bioenergetic or a survival pathway? *Trends Biochem. Sci.* 35, 145–149. doi: 10.1016/j.tibs.2009.10.006
49. Pellerin, L., and Magistretti, P. J. (1994). Glutamate uptake into astrocytes stimulates aerobic glycolysis: a mechanism coupling neuronal activity to glucose utilization. *Proc. Natl. Acad. Sci. U.S.A.* 91, 10625–10629. doi: 10.1073/pnas.91.22.10625
50. Pellerin, L., Pellegrini, G., Bittar, P. G., Charnay, Y., Bouras, C., Martin, J. L., et al. (1998). Evidence supporting the existence of an activity-dependent astrocyte-neuron lactate shuttle. *Dev. Neurosci.* 20, 291–299. doi: 10.1159/000017324
51. Allaman, I., Bélanger, M., and Magistretti, P. J. (2011). Astrocyte-neuron metabolic relationships: for better and for worse. *Trends Neurosci.* 34, 76–87. doi: 10.1016/j.tins.2010.12.001
52. Barros, L. F., Courjaret, R., Jakoby, P., Loaiza, A., Lohr, C., and Deitmer, J. W. (2009). Preferential transport and metabolism of glucose in Bergmann glia over Purkinje cells: a multiphoton study of cerebellar slices. *Glia* 57, 962–970. doi: 10.1002/glia.20820
53. Chuquet, J., Quilichini, P., Nimchinsky, E. A., and Buzsáki, G. (2010). Predominant enhancement of glucose uptake in astrocytes versus neurons during activation of the somatosensory cortex. *J. Neurosci.* 30, 15298–15303. doi: 10.1523/JNEUROSCI.0762-10.2010
54. Casson, R. J., Wood, J. P. M., Han, G., Kittipassorn, T., Peet, D. J., and Chidlow, G. (2016). M-Type pyruvate kinase isoforms and lactate dehydrogenase A in the mammalian retina: metabolic implications. *Invest. Ophthalmol. Vis. Sci.* 57, 66–80. doi: 10.1167/iovs.15-17962
55. Pellerin, L., Bergersen, L. H., Halestrap, A. P., and Pierre, K. (2005). Cellular and subcellular distribution of monocarboxylate transporters in cultured brain cells and in the adult brain. *J. Neurosci. Res.* 79, 55–64. doi: 10.1002/jnr.20307
56. Magistretti, P. J. (2006). Neuron-glia metabolic coupling and plasticity. *J. Exp. Biol.* 209, 2304–2311. doi: 10.1242/jeb.02208
57. Tang, B. L. (2018). Brain activity-induced neuronal glucose uptake/glycolysis: is the lactate shuttle not required? *Brain Res. Bull.* 137, 225–228. doi: 10.1016/j.brainresbull.2017.12.010
58. van Hall, G., Strømstad, M., Rasmussen, P., Jans, O., Zaar, M., Gam, C., et al. (2009). Blood lactate is an important energy source for the human brain. *J. Cereb. Blood Flow Metab.* 29, 1121–1129. doi: 10.1038/jcbfm.2009.35
59. Wyss, M. T., Jolivet, R., Buck, A., Magistretti, P. J., and Weber, B. (2011). In vivo evidence for lactate as a neuronal energy source. *J. Neurosci.* 31, 7477–7485. doi: 10.1523/JNEUROSCI.0415-11.2011
60. Ruminot, I., Schmälzle, J., Leyton, B., Barros, L. F., and Deitmer, J. W. (2019). Tight coupling of astrocyte energy metabolism to synaptic activity revealed by genetically encoded FRET nanosensors in hippocampal tissue. *J. Cereb. Blood Flow Metab.* 39, 513–523. doi: 10.1177/0271678X17737012
61. Volkenhoff, A., Weiler, A., Letzel, M., Stehling, M., Klämbt, C., and Schirmeier, S. (2015). Glial glycolysis is essential for neuronal survival in *Drosophila*. *Cell Metab.* 22, 437–447. doi: 10.1016/j.cmet.2015.07.006
62. Supplie, L. M., Düking, T., Campbell, G., Diaz, F., Moraes, C. T., Götz, M., et al. (2017). Respiration-Deficient astrocytes survive as glycolytic cells in vivo. *J. Neurosci.* 37, 4231–4242. doi: 10.1523/JNEUROSCI.0756-16.2017
63. Mazuel, L., Blanc, J., Repond, C., Bouchaud, V., Raffard, G., Déglon, N., et al. (2017). A neuronal MCT2 knockdown in the rat somatosensory cortex reduces both the NMR lactate signal and the BOLD response during whisker stimulation. *PLoS One* 12:e0174990. doi: 10.1371/journal.pone.0174990

64. Mächler, P., Wyss, M. T., Elsayed, M., Stobart, J., Gutierrez, R., von Faber-Castell, A., et al. (2016). In vivo evidence for a lactate gradient from astrocytes to neurons. *Cell Metab.* 23, 94–102. doi: 10.1016/j.cmet.2015.10.010
65. Mongeon, R., Venkatachalam, V., and Yellen, G. (2016). Cytosolic NADH-NAD(+) redox visualized in brain slices by two-photon fluorescence lifetime biosensor imaging. *Antioxid. Redox Signal.* 25, 553–563. doi: 10.1089/ars.2015.6593
66. Jakoby, P., Schmidt, E., Ruminot, I., Gutiérrez, R., Barros, L. F., and Deitmer, J. W. (2014). Higher transport and metabolism of glucose in astrocytes compared with neurons: a multiphoton study of hippocampal and cerebellar tissue slices. *Cereb. Cortex* 24, 222–231. doi: 10.1093/cercor/bhs309
67. Fünfschilling, U., Supplie, L. M., Mahad, D., Boretius, S., Saab, A. S., Edgar, J., et al. (2012). Glycolytic oligodendrocytes maintain myelin and long-term axonal integrity. *Nature* 485, 517–521. doi: 10.1038/nature11007
68. Zimmer, E. R., Parent, M. J., Souza, D. G., Leuzy, A., Lecrux, C., Kim, H.-I., et al. (2017). 18FFDG PET signal is driven by astroglial glutamate transport. *Nat. Neurosci.* 20, 393–395. doi: 10.1038/nn.4492
69. Dowling, J. E. (1970). Organization of vertebrate retinas. *Invest. Ophthalmol.* 9, 655–680.
69. Baden, T., Euler, T., and Berens, P. (2020). Understanding the retinal basis of vision across species. *Nat. Rev. Neurosci.* 21, 5–20. doi: 10.1038/s41583-019-0242-1
70. Bok, D. (1993). The retinal pigment epithelium: a versatile partner in vision. *J. Cell Sci. Suppl.* 17, 189–195. doi: 10.1242/jcs.1993.supplement_17.27
71. Campbell, M., and Humphries, P. (2012). The blood-retina barrier: tight junctions and barrier modulation. *Adv. Exp. Med. Biol.* 763, 70–84.
72. Country, M. W. (2017). Retinal metabolism: a comparative look at energetics in the retina. *Brain Res.* 1672, 50–57. doi: 10.1016/j.brainres.2017.07.025
72. Wong-Riley, M. T. T. (2010). Energy metabolism of the visual system. *Eye Brain* 2, 99–116. doi: 10.2147/EB.S9078
73. Warburg, O., Posener, K., and Negelein, E. (1924). Ueber den Stoffwechsel der Tumoren. *Biochem. Z.* 152, 309–344.
74. Warburg, O. (1956). On the origin of cancer cells. *Science* 123, 309–314. doi: 10.1126/science.123.3191.309
75. Ng, S. K., Wood, J. P. M., Chidlow, G., Han, G., Kittipassorn, T., Peet, D. J., et al. (2015). Cancer-like metabolism of the mammalian retina. *Clin. Exp. Ophthalmol.* 43, 367–376. doi: 10.1111/ceo.12462
76. DeBerardinis, R. J., and Chandel, N. S. (2020). We need to talk about the Warburg effect. *Nat. Metab.* 2, 127–129. doi: 10.1038/s42255-020-0172-2
77. Young, R. W. (1967). The renewal of photoreceptor cell outer segments. *J. Cell Biol.* 33, 61–72. doi: 10.1083/jcb.33.1.61
78. Young, R. W. (1971). The renewal of rod and cone outer segments in the rhesus monkey. *J. Cell Biol.* 49, 303–318. doi: 10.1083/jcb.49.2.303
79. Wang, W., Fernandez de Castro, J., Vukmanic, E., Zhou, L., Emery, D., Demarco, P. J., et al. (2011). Selective rod degeneration and partial cone inactivation characterize an iodoacetic acid model of Swine retinal degeneration. *Invest. Ophthalmol. Vis. Sci.* 52, 7917–7923. doi: 10.1167/iovs.11-7849
80. Wang, W., Lee, S. J., Scott, P. A., Lu, X., Emery, D., Liu, Y., et al. (2016). Two-Step reactivation of dormant cones in retinitis pigmentosa. *Cell Rep.* 15, 372–385. doi: 10.1016/j.celrep.2016.03.022
81. Park, K. S., Xu, C. L., Cui, X., and Tsang, S. H. (2018). Reprogramming the metabolome rescues retinal degeneration. *Cell. Mol. Life Sci.* 75, 1559–1566. doi: 10.1007/s00018-018-2744-9
82. Léveillard, T., Philp, N. J., and Sennlaub, F. (2019). Is retinal metabolic dysfunction at the center of the pathogenesis of age-related macular degeneration? *Int. J. Mol. Sci.* 20, 762. doi: 10.3390/ijms20030762
83. Hsu, S. C., and Molday, R. S. (1991). Glycolytic enzymes and a GLUT-1 glucose transporter in the outer segments of rod and cone photoreceptor cells. *J. Biol. Chem.* 266, 21745–21752.
84. Gospe, S. M., Baker, S. A., and Arshavsky, V. Y. (2010). Facilitative glucose transporter Glut1 is actively excluded from rod outer segments. *J. Cell Sci.* 123, 3639–3644. doi: 10.1242/jcs.072389
85. Veys, K., Fan, Z., Ghobrial, M., Bouché, A., García-Caballero, M., Vriens, K., et al. (2020). Role of the GLUT1 glucose transporter in postnatal CNS angiogenesis and blood-brain barrier integrity. *Circ. Res.* 127, 466–482. doi: 10.1161/CIRCRESAHA.119.316463
86. Watanabe, T., Matsushima, S., Okazaki, M., Nagamatsu, S., Hirose, K., Uchimura, H., et

- al. (1996). Localization and ontogeny of GLUT3 expression in the rat retina. *Brain Res. Dev. Brain Res.* 94, 60–66. doi: 10.1016/0165-3806(96)00044-2
87. Holman, G. D. (2020). Structure, function and regulation of mammalian glucose transporters of the SLC2 family. *Pflugers Arch.* 472, 1155–1175. doi: 10.1007/s00424-020-02411-3
 88. Cheeseman, C., and Long, W. (2015). Structure of, and functional insight into the GLUT family of membrane transporters. *Cell Health Cytoskeleton* 7, 167–183. doi: 10.2147/CHC.S60484
 89. Narayan, D. S., Chidlow, G., Wood, J. P., and Casson, R. J. (2017). Glucose
 90. metabolism in mammalian photoreceptor inner and outer segments. *Clin. Exp. Ophthalmol.* 45, 730–741. doi: 10.1111/ceo.12952
 91. Okawa, H., Sampath, A. P., Laughlin, S. B., and Fain, G. L. (2008). ATP consumption by mammalian rod photoreceptors in darkness and in light. *Curr. Biol.* 18, 1917–1921. doi: 10.1016/j.cub.2008.10.029
 92. Hurley, J. B., Lindsay, K. J., and Du, J. (2015). Glucose, lactate, and shuttling of metabolites in vertebrate retinas. *J. Neurosci. Res.* 93, 1079–1092. doi: 10.1002/jnr.23583
 93. Ingram, N. T., Fain, G. L., and Sampath, A. P. (2020). Elevated energy requirement of cone photoreceptors. *Proc. Natl. Acad. Sci. U.S.A.* 117, 19599–19603. doi: 10.1073/pnas.2001776117
 94. Ames, A., Li, Y. Y., Heher, E. C., and Kimble, C. R. (1992). Energy metabolism of rabbit retina as related to function: high cost of Na⁺ transport. *J. Neurosci.* 12, 840–853. doi: 10.1523/JNEUROSCI.12-03-00840.1992
 95. Hurley, J. B., Lindsay, K. J., and Du, J. (2015). Glucose, lactate, and shuttling of metabolites in vertebrate retinas. *J. Neurosci. Res.* 93, 1079–1092. doi: 10.1002/jnr.23583 Kanow, M. A., Giarmarco, M. M., Jankowski, C. Sr., Tsantilas, K., Engel, A. L., Du, J., et al. (2017). Biochemical adaptations of the retina and retinal pigment epithelium support a metabolic ecosystem in the vertebrate eye. *Elife* 6:e28899. doi: 10.7554/eLife.28899
 96. Namba, T., Nardelli, J., Gressens, P., and Huttner, W. B. (2020). Metabolic regulation of neocortical expansion in development and evolution. *Neuron* 109, 408–419. doi: 10.1016/j.neuron.2020.11.014
 97. Casson, R. J., Wood, J. P. M., Han, G., Kittipassorn, T., Peet, D. J., and Chidlow, G. (2016). M-Type pyruvate kinase isoforms and lactate dehydrogenase A in the mammalian retina: metabolic implications. *Invest. Ophthalmol. Vis. Sci.* 57, 66–80. doi: 10.1167/iovs.15-17962
 98. Badr, G. A., Tang, J., Ismail-Beigi, F., and Kern, T. S. (2000). Diabetes downregulates GLUT1 expression in the retina and its microvessels but not in the cerebral cortex or its microvessels. *Diabetes* 49, 1016–1021. doi: 10.2337/diabetes.49.6.1016
 99. Bisbach, C. M., Hass, D. T., Robbins, B. M., Rountree, A. M., Sadilek, M., Sweet, I. R., et al. (2020). Succinate can shuttle reducing power from the hypoxic retina to the O₂-rich pigment epithelium. *Cell Rep.* 31:107606. doi: 10.1016/j.celrep.2020.107606
 100. Adler, L., Chen, C., and Koutalos, Y. (2014). Mitochondria contribute to NADPH generation in mouse rod photoreceptors. *J. Biol. Chem.* 289, 1519–1528. doi: 10.1074/jbc.M113.511295
 101. Adjianto, J., Du, J., Moffat, C., Seifert, E. L., Hurle, J. B., and Philp, N. J. (2014). The retinal pigment epithelium utilizes fatty acids for ketogenesis. *J. Biol. Chem.* 289, 20570–20582. doi: 10.1074/jbc.M114.565457
 102. Fu, Z., Kern, T. S., Hellström, A., and Smith, L. E. (2021). Fatty acid oxidation and photoreceptor metabolic needs. *J. Lipid Res.* 62:100035. doi: 10.1194/jlr.TR120000618
 103. Reyes-Reveles, J., Dhingra, A., Alexander, D., Bragin, A., Philp, N. J., and Boesze-Battaglia, K. (2017). Phagocytosis-dependent ketogenesis in retinal pigment epithelium. *J. Biol. Chem.* 292, 8038–8047. doi: 10.1074/jbc.M116.770784
 104. Lévillard, T., and Sahel, J.-A. (2017). Metabolic and redox signaling in the retina. *Cell. Mol. Life Sci.* 74, 3649–3665. doi: 10.1007/s00018-016-2318-7 Wang, W., Kini, A., Wang, Y., Liu, T., Chen, Y., Vukmanic, E., et al. (2019). Metabolic deregulation of the blood-outer retinal barrier in retinitis pigmentosa. *Cell Rep.* 28, 1323–1334.e4. doi: 10.1016/j.celrep.2019.06.093
 105. Petit, L., Ma, S., Cipi, J., Cheng, S.-Y., Zieger, M., Hay, N., et al. (2018). Aerobic glycolysis is essential for normal rod function and controls secondary cone death in retinitis pigmentosa. *Cell Rep.* 23, 2629–2642. doi: 10.1016/j.celrep.2018.04.111
 106. Chinchore, Y., Begaj, T., Wu, D., Drokhlyansky, E., and Cepko, C. L. (2017). Glycolytic reliance promotes anabolism in photoreceptors. *Elife* 6:e25946. doi: 10.7554/eLife.25946

107. Ripps, H. (2002). Cell death in retinitis pigmentosa: gap junctions and the 'bystander' effect. *Exp. Eye Res.* 74, 327–336. doi: 10.1006/exer.2002.1155
108. L  veillard, T., Mohand-S  id, S., Lorentz, O., Hicks, D., Fintz, A.-C., Cl  rin, E., et al. (2004). Identification and characterization of rod-derived cone viability factor. *Nat. Genet.* 36, 755–759. doi: 10.1038/ng1386
109. A  t-Ali, N., Fridlich, R., Millet-Puel, G., Cl  rin, E., Delalande, F., Jaillard, C., et al. (2015). Rod-derived cone viability factor promotes cone survival by stimulating aerobic glycolysis. *Cell* 161, 817–832. doi:10.1016/j.cell.2015.03.023
110. Giarmarco, M. M., Brock, D. C., Robbins, B. M., Cleghorn, W. M., Tsantilas, K. A., Kuch, K. C., et al. (2020). Daily mitochondrial dynamics in cone photoreceptors. *Proc. Natl. Acad. Sci. U.S.A.* 117, 28816–28827. doi: 10.1073/pnas.2007827117
111. Campochiaro, P. A., and Mir, T. A. (2018). The mechanism of cone cell death in Retinitis Pigmentosa. *Prog. Retin. Eye Res.* 62, 24–37. doi: 10.1016/j.preteyeres.2017.08.004
112. Mitchell, P., Liew, G., Gopinath, B., and Wong, T. Y. (2018). Age-related macular degeneration. *Lancet* 392, 1147–1159. doi: 10.1016/S0140-6736(18)31550-2
113. Ferrington, D. A., Ebeling, M. C., Kapphahn, R. J., Terluk, M. R., Fisher, C. R., Polanco, J. R., et al. (2017). Altered bioenergetics and enhanced resistance to oxidative stress in human retinal pigment epithelial cells from donors with age-related macular degeneration. *Redox Biol.* 13, 255–265. doi: 10.1016/j.redox.2017.05.015
114. Hwang, N., Kwon, M.-Y., Woo, J. M., and Chung, S. W. (2019). Oxidative stress-induced pentraxin 3 expression in human retinal pigment epithelial cells is involved in the pathogenesis of age-related macular degeneration. *Int. J. Mol. Sci.* 20:6028. doi: 10.3390/ijms20236028
115. Terluk, M. R., Ebeling, M. C., Fisher, C. R., Kapphahn, R. J., Yuan, C., Kartha, R. V., et al. (2019). N-Acetyl-L-cysteine protects human retinal pigment epithelial cells from oxidative damage: implications for age-related macular degeneration. *Oxid. Med. Cell. Longev.* 2019:5174957. doi:10.1155/2019/5174957
116. Terluk, M. R., Kapphahn, R. J., Soukup, L. M., Gong, H., Gallardo, C., Montezuma, S. R., et al. (2015). Investigating mitochondria as a target for treating age-related macular degeneration. *J. Neurosci.* 35, 7304–7311. doi: 10.1523/JNEUROSCI.0190-15.2015
117. Ebeling, M. C., Polanco, J. R., Qu, J., Tu, C., Montezuma, S. R., and Ferrington, D. A. (2020). Improving retinal mitochondrial function as a treatment for age-related macular degeneration. *Redox Biol.* 34:101552. doi: 10.1016/j.redox.2020.101552
118. Karunadharma, P. P., Nordgaard, C. L., Olsen, T. W., and Ferrington, D. A. (2010). Mitochondrial DNA damage as a potential mechanism for age-related macular degeneration. *Invest. Ophthalmol. Vis. Sci.* 51, 5470–5479. doi: 10.1167/iovs.10-5429
119. Hartong, D. T., Berson, E. L., and Dryja, T. P. (2006). Retinitis pigmentosa. *Lancet* 368, 1795–1809. doi: 10.1016/S0140-6736(06)69740-7
120. Bhandari SK, Lee S, Kim H. Integrative Landscape of Dry AMD Pathogenesis, Models, and Emerging Therapeutic Strategies. *Int J Mol Sci* 2025;27(1):202. doi: 10.3390/ijms27010202.
121. Hansman DS, Jianhai Du b, Casson-RJ, c, Peet DJ. Eye on the horizon: The metabolic role of the RPE in aging and disease. *Prog. Retinal and Eye Research* 2025;104:1-6 a
122. Kulvinder Kochar Kaur, Allahbadia GN, Singh M. The importance of Dysbiosis in Intestinal flora Subsequent to ischaemic Stroke: Implications in Therapeutic Management and Biomarkers for Prognosis-A Narrative Review, *J. Clinical and Medical Research and Studies.* 2023; 2(1):1-13.
123. Kulvinder Kochar Kaur, Allahbadia GN, Singh M. An Update on Role of Bile Acids in Neurological Functions and Neurodegenerative Diseases: A Narrative Review. *J Clin Biomed Invest.* 2023;3(2): 22-39. doi: 10.52916/jcbi224026
124. Kulvinder Kochar Kaur, Allahbadia GN, Singh M. A Comprehensive Update on the Etiopathogenesis of Amyotrophic Lateral Sclerosis with Specific Emphasis on Gut Microbiota (GM), Enteric Nervous System (ENS) and Associated Crosstalk of Astrocytes, GM, Muscle with Mitochondrial Melatonergic Pathway-A Narrative Review. *Int J Neurobiol* 2023; 5(2): 158 DOI: <https://doi.org/10.36266/IJN/158>.
125. Kulvinder Kochar Kaur, Allahbadia GN, Singh M. An Update on Mitophagy Facilitating Substances: Their Future for Facilitating Healthy Ageing Along with Treating NDG (Parkinson's Disease; Alzheimer's Disease), Cancer, Ovarian Ageing: A Mini Narrative Review. *British Journal of Healthcare and Medical Research, Vol* 2024;- 11(5). 28-47. DOI:10.14738/bjhm:115.17586

126. Kulvinder Kochar Kaur, Allahbadia GN, Singh M'. Combined use of Cell Death Mechanisms as plausible therapeutic targets with canonical therapies for Breast Cancer: potentially decreasing dose, toxicity, expenditure & improved survival -A narrative review". Accepted for Publication in Women Health Care & Gynaecology 2025.
127. Kulvinder Kochar Kaur, Gautam Nand Allahbadia, Mandeep Singh. How to Attain Clinical Translation Using Interactions Amongst Endoplasmic Reticulum Stress and Ferroptosis as Therapeutic targets for Improvement of Outcomes & Prognosis of Ovarian Cancers: A Comprehensive Narrative Review, J Clinical and Medical Research and Studies, 2025; V(4)I(5):1-14, DOI: 10.59468/2836-8525/126
128. Kulvinder Kochar Kaur, Allahbadia GN. An update on Targeting Ferroptosis for generating innovative strategies in treatment of Diabetic Kidney Disease: a narrative review" Med Discoveries 2025; 4(6): 1262
129. Kulvinder Kochar Kaur, Gautam Nand Allahbadia, Mandeep Singh An Update on the Manner Adenylyl Cycles (ACS) Might be the Therapeutic Targets Regarding Neurodegenerative Diseases (NDD) Treatment: A Review J. of Adv Clin Neu Res 2025.; 1(2), 01-29. WMJ/JACNR-110 DOI: doi.org/10.63721/25JACNR0110.
130. Kulvinder Kochar Kaur, Allahbadia GN, Singh M. "Targeting Mechanistically interaction Amongst neuron-glia redox signalling subsequent to CNS damage/neurodegenerative diseases (NDD) generation: specifically Astrocytic antioxidant mechanistic modes -A Review" J of Adv Clin Neu Re 2026; 2(1), 01-34. WMJ/JACNR-114 DOI: doi.org/10.63721/26JACNR0114
131. Lee, H., Yoon, D.E., Kim, K., 2020. Genome editing methods in animal models. Anim. Cell Syst. 24, 8–16. <https://doi.org/10.1080/19768354.2020.1726462>.
132. Pennesi, M.E., Neuringer, M., Courtney, R.J., 2012. Molecular Aspects of Medicine Animal models of age related macular degeneration. Mol. Aspect. Med. 33, 487–509. <https://doi.org/10.1016/j.mam.2012.06.003>
133. Petit, L., Ma, S., Cipi, J., Cheng, S.Y., Zieger, M., Hay, N., Punzo, C., 2018. Aerobic glycolysis is essential for normal rod function and controls secondary cone death in retinitis pigmentosa. Cell Rep. 23, 2629–2642. <https://doi.org/10.1016/j.celrep.2018.04.111>.
134. Zhou, Z., Doggett, T.A., Sene, A., Apte, R.S., Ferguson, T.A., 2015. Autophagy supports survival and phototransduction protein levels in rod photoreceptors. Cell Death Differ. 22, 488–498. <https://doi.org/10.1038/cdd.2014.229>.
135. Cheng, S.Y., Cipi, J., Ma, S., Hafler, B.P., Kanadia, R.N., Brush, R.S., Agbaga, M.P., Punzo, C., 2020. Altered photoreceptor metabolism in mouse causes late stage age-related macular degeneration-like pathologies. Proc. Natl. Acad. Sci. U.S.A. 117, 13094–13104. <https://doi.org/10.1073/pnas.2000339117>
136. Dong, A., Xie, B., Shen, J., Yoshida, T., Yokoi, K., Hackett, S.F., Campochiaro, P.A., 2009. Oxidative stress promotes ocular neovascularization. J. Cell. Physiol. 219, 544–552. <https://doi.org/10.1002/jcp.21698>.
137. Chuang, J.Z., Yang, N., Nakajima, N., Otsu, W., Fu, C., Yang, H.H., Lee, M.P., Akbar, A.F., Badea, T.C., Guo, Z., Nuzuzaman, A., Hsu, K.S., Dunaief, J.L., Sung, C.H., 2022. Retinal pigment epithelium-specific CLIC4 mutant is a mouse model of dry age-related macular degeneration. Nat. Commun. 13, e374. <https://doi.org/10.1038/s41467-021-27935-9>
138. Li, K., Wu, J., Lin, Q., Fan, Z., Tang, Q., Lu, W., Zhang, Haipeng, Xing, F., Meng, M., Zou, S., Wei, W., Chen, H., Cai, Jian, Wang, H., Zhang, Hui, Cai, Jiayue, Fang, Ling, Bian, X., Chen, C., Lan, P., Ghesquière, B., Fang, Lekun, Lee, M.H., 2020b. ILF3 is a substrate of SPOP for regulating serine biosynthesis in colorectal cancer. Cell Res. 30, 163–178. <https://doi.org/10.1038/s41422-019-0257-1>
139. Shao, A., Lopez, A.J., Chen, J., Tham, A., Javier, S., Quiroz, A., Frick, S., Levine, E.M., Lloyd, K.C.K., Leonard, B.C., Murphy, C.J., Glaser, T.M., Moshiri, A., 2022. Arap1 loss causes retinal pigment epithelium phagocytic dysfunction and subsequent photoreceptor death. Dis. Model. Mech. 15, dmm049343. <https://doi.org/10.1242/dmm.049343>.
140. Zhang, J., Song, F., Zhao, X., Jiang, H., Wu, X., Wang, B., Zhou, M., Tian, M., Shi, B., Wang, Huamao, Jia, Y., Wang, Hai, Pan, X., Li, Z., 2017a. EGFR modulates monounsaturated fatty acid synthesis through phosphorylation of SCD1 in lung cancer. Mol. Cancer 16, 127. <https://doi.org/10.1186/s12943-017-0704-x>.
141. Sinha, T., Ikelle, L., Naash, M.I., Al-Ubaidi, M.R., 2020a. The intersection of serine metabolism and cellular dysfunction in retinal degeneration. Cells 9, 674. <https://doi.org/10.3390/cells903067>

142. Zhang, R., Engel, A.L., Wang, Y., Li, B., Shen, W., Gillies, M.C., Chao, J.R., 2021. Inhibition of mitochondrial respiration impairs nutrient consumption and metabolite transport in human retinal pigment epithelium. *J. Proteome Res.* 20, 909–922. <https://doi.org/10.1021/acs.jproteome.0c00690>.
143. Zhang, Y., Cross, S.D., Stanton, J.B., Marmorstein, A.D., Le, Y.Z., Marmorstein, L.Y., 2017b. Early AMD-like defects in the RPE and retinal degeneration in aged mice with RPE-specific deletion of Atg5 or Atg7. *Mol. Vis.* 23, 228–241.
144. Chao, J.R., Knight, K., Engel, A.L., Jankowski, C., Wang, Y., Manson, M.A., Gu, H., Djukovic, D., Raftery, D., Hurley, J.B., Du, J., 2017. Human retinal pigment epithelial cells prefer proline as a nutrient and transport metabolic intermediates to the retinal side. *J. Biol. Chem.* 292, 12895–12905. <https://doi.org/10.1074/jbc.M117.788422>.
145. Du, J., Yanagida, A., Knight, K., Engel, A.L., Vo, A.H., Jankowski, C., Sadilek, M., Tran, V. T.B., Manson, M.A., Ramakrishnan, A., Hurley, J.B., Chao, J.R., 2016. Reductive carboxylation is a major metabolic pathway in the retinal pigment epithelium. *Proc. Natl. Acad. Sci. U.S.A.* 113, 14710–14715. <https://doi.org/10.1073/pnas.1604572113>.
146. Yam, M., Engel, A.L., Wang, Y., Zhu, S., Hauer, A., Zhang, R., Lohner, D., Huang, J., Dinterman, M., Zhao, C., Chao, J.R., Du, X.J., 2019. Proline mediates metabolic communication between retinal pigment epithelial cells and the retina. *J. Biol. Chem.* 294, 10278–10289. <https://doi.org/10.1074/jbc.RA119.007983>.
147. Hass, D.T., Bisbach, C.M., Robbins, B.M., Sadilek, M., Sweet, I.R., Hurley, J.B., 2022. Succinate metabolism in the retinal pigment epithelium uncouples respiration from ATP synthesis. *Cell Rep.* 39, 110917. <https://doi.org/10.1016/j.celrep.2022.110917>.
148. Gillies, R.J., Morse, D.L., 2005. In vivo magnetic resonance spectroscopy in cancer. *Annu. Rev. Biomed. Eng.* 7, 287–326. <https://doi.org/10.1146/annurev.bioeng.7.060804.100411>.
149. Van Der Graaf, M., 2010. In vivo magnetic resonance spectroscopy: basic methodology and clinical applications. *Eur. Biophys. J.* 39, 527–540. <https://doi.org/10.1007/s00249-009-0517-y>.
150. Buonocore, M.H., Maddock, R.J., 2015. Magnetic resonance spectroscopy of the brain: a review of physical principles and technical methods. *Rev. Neurosci.* 26, 609–632. <https://doi.org/10.1515/revneuro-2015-001>.
151. Vachha, B., Huang, S.Y., 2021. MRI with ultrahigh field strength and high-performance gradients: challenges and opportunities for clinical neuroimaging at 7 T and beyond. *Eur. Radiol. Exp.* 5, 35. <https://doi.org/10.1186/s41747-021-00216-2>.
152. Bongaerts, J., De Bundel, D., Mangelings, D., Smolders, I., Vander Heyden, Y., Van Eeckhaut, A., 2018. Sensitive targeted methods for brain metabolomic studies in microdialysis samples. *J. Pharm. Biomed. Anal.* 161, 192–205. <https://doi.org/10.1016/j.jpba.2018.08.043>.
153. Kao, C.-Y., Anderzhanova, E., Asara, J.M., Wotjak, C.T., Turck, C.W., 2015. NextGen brain microdialysis: applying modern metabolomics technology to the analysis of extracellular fluid in the central nervous system. *Complex Psychiatry* 1, 60–67. <https://doi.org/10.1159/000381855>.
154. Louzada-Junior, P., Dias, J.J., Santos, W.F., Lachat, J.J., Bradford, H.F., CoutinhoNetto, J., 1992. Glutamate release in experimental ischaemia of the retina: an approach using microdialysis. *J. Neurochem.* 59, 358–363. <https://doi.org/10.1111/j.1471-4159.1992.tb08912.x>.
155. Kataoka, K., Matsumoto, H., Kaneko, H., Notomi, S., Takeuchi, K., Sweigard, J.H., Atik, A., Murakami, Y., Connor, K.M., Terasaki, H., Miller, J.W., Vavvas, D.G., 2015. Macrophage- and RIP3-dependent inflammasome activation exacerbates retinal detachment-induced photoreceptor cell death. *Cell Death Dis.* 6, e1731. <https://doi.org/10.1038/cddis.2015.73>.
156. Gilmore, I.S., Heiles, S., Pieterse, C.L., 2019. Metabolic imaging at the single-cell scale: recent advances in mass spectrometry imaging. *Annu. Rev. Anal. Chem.* 12, 201–224. <https://doi.org/10.1146/annurev-anchem-061318-115516>.
157. Taylor, M.J., Lukowski, J.K., Anderton, C.R., 2021. Spatially resolved mass spectrometry at the single cell: recent innovations in proteomics and metabolomics. *J. Am. Soc. Mass Spectrom.* 32, 872–894. <https://doi.org/10.1021/jasms.0c00439>.
158. Sun, N., Ly, A., Meding, S., Witting, M., Hauck, S.M., Ueffing, M., Schmitt-Kopplin, P., Aichler, M., Walch, A., 2014. High-resolution metabolite imaging of light

- and dark treated retina using MALDI-FTICR mass spectrometry. *Proteomics* 14, 913–923. <https://doi.org/10.1002/pmic.201300407>.
159. Anderson, D.M.G., Ablonczy, Z., Koutalos, Y., Spragins, J., Crouch, R.K., Caprioli, R.M., Schey, K.L., 2014. High resolution MALDI imaging mass spectrometry of retinal tissue lipids. *J. Am. Soc. Mass Spectrom.* 25, 1394–1403. <https://doi.org/10.1007/s13361-014-0883-2>.

Cite this article: Kulvinder Kochar Kaur, *Further Expanding the Metabolic Topography for Sustenance of Completeness Along with Working of Retina –A Review International Journal of Research in Medical and Clinical Science (IJRMCS)* Online ISSN:2997-6189 2026; 4(1): 209-230.

Copyright: © 2026. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.